

## Learning Genetics Can Be Fun - Solutions

1. Two black dogs could be homozygous black (BB) or heterozygous black (Bb). Yellow must be homozygous, therefore cannot be the same genotype as black.

2. P Cc x Cc            both parents are normal but "carry" the allele for CF. One in four children will  
F<sub>1</sub> CC, Cc, Cc, cc    inherit it.

3. a) P Rr x rr  
b) R, r and r, r  
F<sub>1</sub> Rr, Rr, rr, rr        1 round:1 wrinkled  
F<sub>2</sub> RR, Rr, Rr, rr       3 round:1 wrinkled

4. P Ll x ll  
F<sub>1</sub> Ll, Ll, ll, ll            1 long:1 short

5. P T<sub>-</sub> x tt                - almost 1:1 therefore unknown parent must be heterozygous.  
F<sub>1</sub> 327 tall: 321 short Note: homozygous (TT) would give ALL tall plants in F<sub>1</sub>.

6. The presence of all smooth in the offspring means smooth is dominant.

P SS x ss  
F<sub>1</sub> Ss  
F<sub>2</sub> 3:1

7. a) P Ss x Ss  
F<sub>1</sub> SS, Ss, Ss, ss  
b) P S<sub>-</sub> x <sub>-</sub>

The female must be heterozygous as she produced non-spotted pups. The unknown male must be homozygous recessive (ss). If he were homozygous dominant, all pups would be spotted. If he were heterozygous, you would expect a 3:1 ratio in pups.

8. (i) P T<sub>-</sub> x tt                The male must be heterozygous (Tt) to be able to produce  
F<sub>1</sub> tt                            both trotters and pacers. If he were homozygous dominant  
(ii) P T<sub>-</sub> x tt                he would produce only trotters.  
F<sub>1</sub> Tt  
(iii) P T<sub>-</sub> x T<sub>t</sub>  
F<sub>1</sub> tt

9. Normal woman Pp (must be heterozygous because father was albino)

Husband pp  
Husband's parents both Pp  
Children Pp, Pp, pp

10. test cross    W<sub>-</sub> x ww

11. B - black; b - white; S - short; s - long

a) P BBSs x bbss

F<sub>1</sub> BbSs, Bbss                      1 black, short:1 black, long

b) P BbSs x bbss

F<sub>1</sub> BbSs, Bbss, bbSs, bbss      1 black, short:1 black, long: 1 white, short:1 white, long

c) P BBss x BbSs

F<sub>1</sub> BBSs, BBss, BbSs, Bbss      1 black, short:1 black, long

d) i) (a) 1/2      (b) 1/4      (c) 1/2

ii) (a) 1/2      (b) 1/4      (c) 1/2

iii) (a) 0      (b) 1/4      (c) 0

12. B - black; b - white; S - solid; s - spotted

                    male                      female

a) P    B\_S\_ x            bbS\_

F<sub>1</sub>      2 BbS\_ , 2 bbS\_

Some white pups so the male must be Bb. The absence of any non-spotted pups suggests that female A is SS but we can't say for sure.

b) P    BbS\_ x            B\_S\_

F<sub>1</sub>      bbss    the presence of white, non-spotted pups means that female B must be BbSs

c) P    BbSs x            bbss

F<sub>1</sub>      bbS\_ , bbss , BbS\_ , Bbss

The genotype of female C can be determined from her phenotype.

13. P Pp x Pp

F<sub>1</sub> PP, Pp, Pp, pp            chance of PKU is 1/4

14.    P Bb x Bb

F<sub>1</sub> BB, Bb, Bb, bb

a) 1/4              (b) 1/4              (c) 1/2

d) 1 homozygous brown:2 heterozygous brown:1 homozygous blonde. Phenotype: 3 brown:1 blonde

e) not possible because blonde (b) is recessive

f) C = curly; c = straight

g) P Cc x cc

F<sub>1</sub> Cc, Cc, cc, cc

h) C, c              (i) c, c              (j) 0              (k) 1/2              (l) 1/2

m) 1 heterozygous:1 homozygous recessive. Phenotype: 1 curly:1 straight

n) No. Straight hair is recessive so individual MUST be homozygous (cc).

15. S<sup>R</sup> - round; S<sup>L</sup> - long

P S<sup>R</sup>S<sup>R</sup> x S<sup>L</sup>S<sup>L</sup>

F<sub>1</sub> S<sup>R</sup>S<sup>L</sup>, S<sup>R</sup>S<sup>L</sup>, S<sup>R</sup>S<sup>L</sup>, S<sup>R</sup>S<sup>L</sup>

P S<sup>R</sup>S<sup>L</sup> x S<sup>R</sup>S<sup>L</sup>

F<sub>2</sub> S<sup>R</sup>S<sup>R</sup>, S<sup>R</sup>S<sup>L</sup>, S<sup>R</sup>S<sup>L</sup>, S<sup>L</sup>S<sup>L</sup>      (incomplete dominance)

16. P S<sup>N</sup>S<sup>M</sup> x S<sup>N</sup>S<sup>M</sup>

$F_1 S^N S^N, S^N S^M, S^M S^M$  25% chance of having homozygous recessive child

17.  $C^R C^R$  - chestnut;  $C^M C^M$  - cremello;  $C^M C^R$  - palomino

P  $C^M C^R \times C^M C^M$

$F_1 C^M C^M, C^R C^M$  1 cremello:1 palomino

18. P  $F^R F^W \times F^R F^W$

$F_1 F^R F^R, F^R F^W, F^R F^W, F^W F^W$

- a)  $\frac{1}{2}$  pink
- b)  $\frac{1}{4}$  red
- c)  $\frac{1}{4}$  white
- d) 1:2:1

19. woman  $I^B \_$  x man  $I^A \_$

$F_1$  ii is possible if mother and father were both heterozygous. The facts are inconclusive.

20. P  $\sigma$  ii x  $\varphi$   $I^A I^B$

$F_1 I^A i, I^B i$

$AB \varphi$  could produce AB offspring if  $\sigma$  were type A, B, or AB; she could never produce type O in  $F_1$  because she always donates either A or B.

21. a) P  $C^h C^a \times C^a C^a$

$F_1 C^h C^a, C^a C^a$  1 himalayan:1 albino

b) P  $CC^a \times C^{ch} C^a$

$F_1 2 C \_, C^{ch} \_, C^a C^a$

c) P  $C^{ch} C^{ch} \times C^{ch} C^a$

$F_1 C^{ch} C^{ch}, C^{ch} C^a$  1 chinchilla:1 light gray

d) P  $C^{ch} \underline{C^a} \times C^a C^a$

$F_1 5 C^h \underline{C^a}, 5 C^{ch} \underline{C^a}$  test cross

22. a) 4 children

b) A is Dd, B is Dd

c) M is dd, N is dd

23.

